

Age ▶ Vaccine	Birth	2 month	4 month	6 month	9 month	12 month	15 month	18 month	2 year	4-6 year	11-18year
Tetanus											Tdap Given at 11-12 years
Diphtheria, Tetanus, Pertussis – DTaP		Pediarix	Pediarix	Pediarix			DTaP			DTaP	
Inactivated Poliovirus		Pediarix	Pediarix	Pediarix						IPV	
Hepatitis B	Hep B	Pediarix	Pediarix	Pediarix							
Haemophilus Influenza B		Hib	Hib	Hib		Hib					
Pneumococcal		PCV13	PCV13	PCV13		PCV13	<i>PCV13 if patient has had only PCV7 in past</i>				
							<i>PPSV23 for High Risk Groups</i>				
Rotavirus		Rota	Rota	Rota							
Hepatitis A						Hep A		Hep A	<i>Catchup if not done</i>		
Measles, Mumps, Rubella						MMR				MMR	
Varicella						Var				Var	<i>Second dose catchup</i>
Influenza				Yearly from 6 months and older; 2 shots in first season if under age 9 and no prior influenza vaccine							
Human Papilloma Virus											< 15 yrs HPV x 2 2 doses 6 months apart >15 yrs HPV x 3 3 doses at 0/2/6months
Meningitis											MCV4 Given at 11-12 years Booster dose five years later Unless first dose at age 16 or after
Meningitis B											MenB x 2 At 16 yrs old 2 doses separated by 6 months