

Vaccine	Age	Birth	2 month	4 month	6 month	9 month	12 month	15 month	18 month	2 year	4 year	11 year	16 year
Diphtheria, Tetanus, Pertussis (DTaP)			Vaxelis	Vaxelis	Vaxelis			DTaP			Kinrix		
Inactivated Poliovirus (IPV)													
Hepatitis B (HepB)													
Haemophilus Influenza B (HIB)							HIB						
Pneumococcal Conjugate (PCV20)			PCV20	PCV20	PCV20		PCV20			PPSV23 for High Risk Groups			
Rotavirus (Rota)			Rota	Rota	Rota								
Hepatitis A (Hep A)							Hep A		Hep A				
Measles, Mumps, Rubella (MMR)							MMR				MMR		
Varicella (VZV)								VZV			VZV		
Influenza					Yearly from 6 months and older; 2 doses in first season if under age 9 and no prior influenza vaccine								
COVID					6mo-4 yrs- See CDC Guidelines- typically 1 or 2 primary doses then 1 seasonal dose							5 years and over - 1 seasonal dose	
Respiratory Syncytial Virus (RSV)	Infants younger than age 8 months who are born shortly before or are entering their first RSV season (typically fall through spring)												
Tetanus, Diptheria, Pertussis (TDaP)												TDaP Given at 11-12 years	
Human Papilloma Virus (HPV9)												HPV9 x2 <15 yrs (9-14 yrs) 2 doses separated by 6 months	HPV9 x3 ≥15 yrs 3 doses given at 0/2/6 months
Meningitis (MenACWY-TT)												MenQuadfi Given at 11-12 years	MenQuadfi Booster approx 4 years after dose 1 (age 16)
Meningitis B (MenB)													MenB x 2 Given at 16 years 2 doses separated by 6 months